



**QUEEN'S
UNIVERSITY
BELFAST**

Metabolic and neuroprotective effects of dapagliflozin and liraglutide in diabetic mice

Millar, P., Pathak, N. M., Parthsarathy, V., Bjourson, A., O'Kane, M., Moffett, R. C., Flatt, P. R., & Pathak, V. (2017). Metabolic and neuroprotective effects of dapagliflozin and liraglutide in diabetic mice. *Journal of Endocrinology*.

Published in:
Journal of Endocrinology

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2017 Society for Endocrinology.

This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Metabolic and neuroprotective effects of dapagliflozin and liraglutide in diabetic mice

Paul Millar¹, Nupur Pathak¹, Vadivel Parthsarathy¹, Anthony Bjourson², Maurice O’Kane^{2,3}, Varun Pathak¹, Charlotte Moffett¹, Peter Flatt¹, Victor Gault¹

¹SAAD Centre for Pharmacy and Diabetes, School of Biomedical Sciences, University of Ulster, Coleraine BT52 1SA, Northern Ireland, UK

²Northern Ireland Centre for Stratified Medicine, University of Ulster, C-TRIC Building, Altnagelvin Hospital, Londonderry BT47 6SB, Northern Ireland, UK

³Clinical Chemistry Laboratory, Western Health and Social Care Trust, Altnagelvin Hospital, Londonderry BT47 6SB, Northern Ireland, UK

Correspondence should be addressed to V Gault
Email: va.gault@ulster.ac.uk

Abstract

This study assessed the metabolic and neuroprotective actions of the sodium glucose co-transporter-2 inhibitor dapagliflozin in combination with the GLP-1 agonist liraglutide in dietary-induced diabetic mice. Mice administered low-dose streptozotocin (STZ) on a high fat diet received dapagliflozin, liraglutide, dapagliflozin-plus-liraglutide (DAPA-Lira) or vehicle once-daily over 28 days. Energy intake, body weight, glucose and insulin concentrations were measured at regular intervals. Glucose tolerance, insulin sensitivity, hormone and biochemical analysis, dual-energy x-ray absorptiometry densitometry, novel object recognition, islet and brain histology were examined. Once-daily administration of DAPA-Lira resulted in significant decreases in body weight, fat mass, glucose and insulin concentrations, despite no change in energy intake. Similar beneficial metabolic improvements were observed regarding glucose tolerance, insulin sensitivity, HOMA-IR, HOMA- β , HbA1c, and triglycerides. Plasma glucagon, GLP-1 and IL-6 levels were increased and corticosterone concentrations decreased. DAPA-Lira treatment decreased alpha cell area and increased insulin content compared to dapagliflozin monotherapy. Recognition memory was significantly improved in all treatment groups. Brain histology demonstrated increased staining for doublecortin (number of immature neurons) in dentate gyrus and synaptophysin (synaptic density) in stratum oriens and stratum pyramidale. These data demonstrate that combination therapy of dapagliflozin and liraglutide exerts beneficial metabolic and neuroprotective effects in diet-induced diabetic mice. Our results highlight important personalised approach in utilising liraglutide in combination with dapagliflozin, instead of either agent alone, for further clinical evaluation in treatment of diabetes and associated neurodegenerative disorders.

Keywords: dapagliflozin; diabetes; GLP-1; glucagon; liraglutide

Introduction

Type 2 diabetes mellitus (T2DM) is a metabolic disorder that arises due to a complex array of molecular defects manifesting in dysregulated insulin secretion, impaired insulin action, or both. Since the pathophysiology of T2DM is multifaceted and involves a range of biochemical mechanisms, there is no single therapy can effectively manage all aspects of the disorder (Zaccardi *et al.* 2016). Moreover, as T2DM and obesity levels are increasing at an alarming rate, more effective therapies and innovative treatment strategies are urgently needed to control glycaemia, reduce body weight and decrease the risk of micro- and macrovascular complications (da Rocha Fernandes *et al.* 2016). The previous two decades have witnessed a surge in the number of new drug classes such as glucagon-like peptide-1 (GLP-1) agonists, dipeptidylpeptidase-4 (DPP4) inhibitors and sodium-glucose-cotransporter-2 (SGLT2) inhibitors (Bailey *et al.* 2016). Although these agents may be used as monotherapies, it is becoming increasingly apparent that successful and cost effective management of T2DM requires development of safe combination therapies with distinct and complementary mechanisms of action.

The kidneys play a pivotal role in regulating glucose homeostasis as most of the glucose filtered by the glomerulus is reabsorbed (Gerich *et al.* 2001). In healthy subjects, the high capacity, low-affinity SGLT2, reabsorbs approximately 90% of glucose in S1 segment of proximal tubules (Hediger & Rhoads 1994, Han *et al.* 2008). Under conditions of chronic hyperglycaemic, SGLT2 is up regulated and this enhances glucose reabsorption and worsens glycaemia (Rahmoune *et al.* 2005). Dapagliflozin, a highly selective and potent oral inhibitor of SGLT2, reduces reabsorption of filtered glucose leading to increased glucosuria and improvement in glycaemic control (Vivian 2015). Although actions of dapagliflozin appear to be independent of insulin secretion, dapagliflozin improves insulin sensitivity, most likely as a result of sustained reduction in hyperglycaemia, alleviation of glucose toxicity and weight reduction through enhanced caloric loss (Macdonald *et al.*

2010, Mudaliar *et al.* 2014, Merovci *et al.* 2015, Millar *et al.* 2016). Beneficial actions of dapagliflozin are, to some extent, limited by unrestrained hepatic glucose production (Bonner *et al.* 2015). Thus, inhibition of hepatic glucose output by stimulation of insulin secretion as well as inhibition of glucagon secretion may significantly enhance therapeutic efficacy of SGLT2 inhibition.

GLP-1 agonists are well established as effective agents to treat patients with T2DM due to a range of beneficial actions including weight loss, induction of satiety, inhibition of gastric emptying, stimulation of insulin secretion and inhibition of alpha cell function (Bailey *et al.* 2016). In addition, GLP-1 agonists exert effects at other extra pancreatic sites (Renner *et al.* 2016), with notable neuroprotective actions in animal models of diabetes-obesity, Alzheimer's disease (AD) and Parkinson's disease (PD) (Ashraghi *et al.* 2016, Tramutola *et al.* 2017). Liraglutide (Victoza®) is a highly effective long-acting GLP-1 agonist that shares 97% sequence homology with human GLP-1 (Knudsen *et al.* 2000). Structural modifications include amino acid substitution of Lys³⁴ with Arg, and addition of lipophilic C₁₆ acyl moiety at position 26 via gamma-glutamyl linker (Madsen *et al.* 2007). These structural changes provide liraglutide with enhanced pharmacokinetic profile and significantly prolonged half-life, thus facilitating once-daily injection (Agersø *et al.* 2002). This prolonged bioactivity has been attributed to non-covalent reversible albumin binding, ability of liraglutide to self-aggregate and form heptamers in solution, and stability to the enzyme DPP4 (Knudsen *et al.* 2000, Madsen *et al.* 2007, Li *et al.* 2016).

Given the need for more personalised treatment strategies for patients with T2DM and the unique mechanism of action of dapagliflozin and liraglutide, we hypothesised that combining both drugs would provide additive metabolic and neuroprotective outcomes. We chose to administer a GLP-1 agonist rather than DPP-4 inhibitor as DPP-4 inhibitors act to prevent degradation of a number of regulatory peptides including GLP-1 (Bailey *et al.* 2016). As such, HF mice on

background low-dose STZ were treated with dapagliflozin or liraglutide as monotherapy and combination therapy for 28 days. Effects on glucose tolerance, insulin sensitivity, body weight, hormones, memory and learning, islet and brain histology were assessed.

Materials and methods

Animals

Male NIH Swiss mice (aged 8-10 weeks) purchased from Harlan (Oxon, UK) were kept at 22±2°C with 12:12 h light/dark cycle. Mice had free access to high fat diet (45% AFE Fat; Product Code 824053; Special Diet Services, Witham, UK; total energy 26.15 kJ/g). An additional group of mice had free access to standard rodent chow (Teklad Global 18% Protein Rodent Diet; Product Code 2018S; Harlan, UK; total energy 13.0 kJ/g). All animals had free access to drinking water and no adverse effects were observed during the entire experimental study. All experiments were performed according to the *Principles of Laboratory Animal Care* (NIH publication no. 86-23, revised 1985) and UK Home Office Regulations (UK Animals Scientific Procedures Act 1986).

Experimental design

Mice commenced high fat diet on day -28 and subsequently received STZ (Sigma-Aldrich, Dorset, UK) prepared in sodium citrate buffer (pH 4.5) on day -14 (50 mg/kg; i.p.) and day -7 (50 mg/kg; i.p.). Mice that displayed a blood glucose concentration greater than 13 mmol/l were recruited into the study. On day 0, mice commenced drug treatments for 28 days as follows: Group 1 (HF control) – high fat mice administered saline vehicle (0.9% wt/vol; p.o.; o.d.); Group 2 (dapagliflozin) – high fat mice administered dapagliflozin (1 mg/kg; p.o.; o.d.; Selleck Chemicals; Stratech Scientific Ltd., Suffolk, UK; Catalog number S1548-SEL); Group 3 (DAPA-Lira) – high fat mice administered dapagliflozin (1 mg/kg; p.o.; o.d.) plus liraglutide (25 nmol/kg; i.p.; o.d.; GL Biochem Ltd.,

Shanghai, China); Group 4 (Lira) - high fat mice administered liraglutide (25 nmol/kg; i.p.; o.d.); Group 5 (lean control) – lean mice administered saline vehicle (0.9% wt/vol; p.o.; o.d.). All treatments administered at 14:00 h and mice remained on respective diet for study duration. The rationale for choosing 1 mg/kg dapagliflozin (p.o.) and 25 nmol/kg liraglutide (i.p.) in this study was based on previously published literature (Moffett *et al.* 2014, Millar *et al.* 2016). Energy intake, body weight, glucose and insulin concentrations were measured every 3 to 4 days. At the end of the study, glucose tolerance (18 mmol/kg; p.o.; at 10:00 h in 12 h-fasted mice), insulin sensitivity (25 U/kg bovine insulin; i.p.; at 10:00 h in non-fasted mice), novel object recognition task, dual-energy x-ray absorptiometry (DEXA) scanning, lipids, hormones/biomarkers, islet and brain histology were performed.

Biochemical and DEXA analyses

Blood samples were collected as indicated in Figures from tail vein of conscious mice into chilled fluoride/heparin micro-centrifuge tubes (Sarstedt, Numbrecht, Germany) and centrifuged at 13,000g for 30 s (Beckman Instruments, Galway, Ireland). Glucose concentrations were measured using Ascencia Contour Blood Glucose Meter (Bayer Healthcare, Newbury, UK) and plasma/pancreatic insulin determined using modified dextran-coated charcoal RIA (Flatt & Bailey 1981). HOMA-IR and HOMA- β were determined from calculations as described previously (Gault *et al.* 2015). Lipids (total-cholesterol – CH200; and triglycerides – TR210) and ALT (AL1205) were measured using enzymatic kits from Randox Laboratories (Crumlin, UK). Plasma corticosterone (ab100712) and IL-6 (ab108821) were measured using enzymatic kits from Abcam (Cambridge, UK) and analysed with SOFTMAX PRO Software Version 5.2 on Flexstation 3 (Molecular Devices, Sunnyvale, CA, USA). Glucagon and total GLP-1 were measured by ELISA (EZGLU-30K and EZGLP1T-36K, respectively; Millipore, UK). HbA1c was determined with a commercially available kit (HB-3058;

Chirus Limited, Watford, UK). Percentage fat and lean mass were measured using DEXA densitometry (Piximus Densitometer, USA) as described previously (Millar *et al.* 2016).

Assessment of learning and memory

Open field and novel recognition tests were performed as described previously (Lennox *et al.* 2014). Briefly, mice were placed in an arena and motor activity (speed and path length), anxiety (grooming events) and exploration (rearing events) recorded over 5 min period. Twenty-four hours later, mice were placed back into the same arena and a novel object recognition task was conducted comprising a 10 min acquisition phase (followed by a 3 h rest in the home cage) followed by test trial where mice could explore familiar and novel object for 10 min. Time spent exploring familiar or novel object was expressed as recognition index (RI) calculated as time (t) spent exploring novel object divided by time spent exploring both objects (A + B) x 10. $RI_B = tB/t(A + B) \times 100$ normalises all data for statistical comparison (Lennox *et al.* 2014).

Immunohistochemistry and image analysis

Mice were perfused with PBS transcardially as described previously (Parthsarathy *et al.* 2013). Pancreatic tissue was excised for immunohistochemistry, measurement of insulin/glucagon content and gene expression. For determination of pancreatic insulin and glucagon content, pre-weighed pancreatic tissue was washed thoroughly in ice-cold PBS, homogenised in acid ethanol solution (ethanol/0.7 M HCl; 3:1 ratio) and extracted overnight at 4°C. Insulin content was measured by insulin radioimmunoassay and glucagon content determined by ELISA (EZGLU-30K; Millipore, UK). For histology, pancreatic tissues were fixed in 4% paraformaldehyde for 48 h at 4°C, processed using automated tissue processor (Leica TP1020, Leica Microsystems, Nussloch, Germany) and embedded in paraffin wax. Immunohistochemistry was performed as described

previously (Moffett *et al.* 2015). Following primary antibodies used: mouse monoclonal anti-insulin antibody (ab6995, 1:1000; Abcam), guinea-pig anti-glucagon antibody (PCA2/4, 1:400; raised in-house), rabbit polyclonal anti-GLP-1 antibody (XJIC8, 1:200; raised in-house) and mouse polyclonal anti-IL-6 details (PM626, 1:200; ThermoFisher Scientific). Secondary antibodies used as appropriate: Alexa Fluor 488 goat anti-guinea pig IgG – 1:400, Alexa Fluor 594 goat anti-mouse IgG – 1:400. Slides were viewed under FITC filter (488 nm) or TRITC filter (594 nm) using fluorescent microscope (Olympus BX51) and DP70 camera adapter system. Brain processing and immunostaining were performed as described previously (Parthasarathy *et al.* 2013). Briefly, 40 micron thick coronal sections of brains at anatomical regions -2 to -3 bregma were stained for young immature neurons (anti-doublecortin, 1:200 dilution, sc-8066, Santa Cruz Biotechnology) and synaptic density (anti-synaptophysin, 1:200 dilution, Abcam, ab-7837).

Image analysis

Alpha and beta cell area were analysed in a blinded manner using Cell^F image analysis software (Olympus Soft Imaging Solutions, GmbH) and expressed as μm^2 . Briefly, fluorescent images were captured using digital camera and closed polygon tool in Cell^F used to analyse alpha cell and beta cell area. Pixel area was converted to μm^2 and plotted in Prism. To quantify cell proliferation and neurogenesis, DCX-labelled immature neurons were counted in sub granular zone of dentate gyrus. Minimum of seven coronal sections per animal were counted using 40 x objective of bright field microscope (Olympus BX51) and plotted as average number of positive cells per section. Synaptophysin staining was analysed with Image J (NIH, USA) software using corrected O.D. method (McClean *et al.* 2011). Briefly, following adjustment for optimum resolution, calibration for optical density was performed using Kodak No. 3 step tablet (Tiffen, Kodak) and calibration curve obtained as described in Image J software. Using 10 x magnification objective, image for each area

of interest was obtained per section (4-5 sections per mouse brain) with digital camera. Area of interest comprised hippocampus and cortex that included polymorphic layer, granular cell layer, molecular layer, stratum radiatum, stratum pyramidal, stratum oriens, interior and exterior cortical layers. Images were converted to 8-bit grey scale and pixel density obtained from three small randomly selected squares per layer converted to O.D. using calibration curve. Average O.D values for each layer were subtracted from average O.D values of granular cell layer (GCL) and corrected O.D. plotted.

Gene expression

mRNA extracted (Tripure Isolation Reagent; Roche Diagnostics, UK), quantified and purity determined using nanophotometer (Implen, Munich, Germany). cDNA synthesized using Transcriptor First Strand cDNA Synthesis Kit (Roche Diagnostics) and gene expression analysis for insulin and glucagon performed on whole pancreas by qPCR using Light Cycler 480 Probes Master (Roche Diagnostics) according to manufacturer's instructions (Gault *et al.* 2015). HPRT and beta-actin were used as internal control for normalisation. PCR conditions were 95°C for 10 min, followed by cDNA amplification for 50 cycles with 95°C denaturation for 10s, 60°C annealing for 30 s and 72°C elongation for 10s followed by cooling period of 30 s at 40 °C. Relative quantification using $2^{-\Delta\Delta CT}$ method used to calculate differences between groups (Livak & Schmittgen 2001).

Statistical analyses

Results were analysed using Prism (GraphPad Software Inc., USA) and data expressed as mean \pm S.E.M. For metabolic data, statistical analyses were performed using one-way ANOVA followed by Student-Newman-Keuls *post-hoc* test. For novel object recognition and immunohistochemistry, statistical analyses were carried out using unpaired Student's t-test (non-parametric, with two-tailed

p values and 95% confidence interval) and one-way ANOVA with Bonferroni *post-hoc* test. Groups of data were considered to be significantly different if $p<0.05$.

Results

Effects of DAPA-Lira on body weight, energy intake, glucose and insulin concentrations

Compared to HF controls, DAPA-Lira treatment resulted in significant time-dependent decrease in body weight ($p<0.001$; Fig. 1A). Importantly, body weights for DAPA-Lira and dapagliflozin groups were reduced despite no reduction in energy intake (Fig. 1B). Liraglutide-treated mice displayed reduced cumulative energy intake ($p<0.05$ - $p<0.001$; Fig. 1B) compared to HF controls. DAPA-Lira treatment resulted in time-dependent decrease (242%; $p<0.001$) in glucose concentrations compared to HF controls, dapagliflozin or liraglutide alone (157-172%; $p<0.01$) on day 28 (Fig. 1C). All treatments exhibited progressive time-dependent increase in insulin but no significance was detected between HF groups, except DAPA-Lira treated mice which exhibited lower levels on day 28 ($p<0.01$; Fig. 1D).

Effects of DAPA-Lira on glucose tolerance, insulin response to glucose, insulin sensitivity, HbA1c and plasma glucagon

Mice treated with DAPA-Lira for 28 days exhibited significant reduction (37-47% decrease; $p<0.01$) in glucose concentrations (individual time-course for up to 120 min) compared with dapagliflozin or liraglutide alone (Fig. 2A). This was further corroborated by significantly reduced glucose AUC₁₂₀ values (37-52% decrease; $p<0.001$; Fig. 2B). As shown in Fig. 2C, all treatment groups demonstrated increased insulintropic response, with DAPA-Lira mice exhibiting significantly higher AUC₉₀ values (1.1-1.4-fold increase; $p<0.01$ - $p<0.001$) compared to dapagliflozin or liraglutide alone (Fig. 2D). Similarly, all treatment groups displayed marked improvement in insulin

sensitivity compared to HF controls following administration of exogenous insulin ($p<0.05$; Fig. 3A-B). DAPA-Lira treated mice also displayed marked reduction (73% lower; $p<0.05$) in HOMA-IR compared to dapagliflozin or liraglutide alone (Fig. 3C). Furthermore, mice treated with DAPA-Lira had a significantly improved HOMA- β index compared to dapagliflozin (53% increase; $p<0.01$) or liraglutide (13% increase; $p<0.05$) alone (Fig. 3D). All treatment groups had significantly ($p<0.001$) reduced HbA1c values compared to HF controls with DAPA-Lira treated mice exhibiting improved HbA1c (19-26% reduction; $p<0.001$) compared to dapagliflozin or liraglutide alone (Fig. 3E). Dapagliflozin and DAPA-Lira groups had significantly increased plasma glucagon concentrations (24-33%; $p<0.001$) compared to HF controls, whereas liraglutide group exerted a 33% reduction in plasma glucagon ($p<0.001$) compared to DAPA-Lira treated mice (Fig. 3F).

Effects of DAPA-Lira on body composition and lipids

DEXA analysis revealed that all treatment groups exhibited significant reduction (37-42% decrease; $p<0.05$ - $p<0.001$) in percentage fat mass compared to HF controls (Fig. 4A). No significant differences were noted between DAPA-Lira and liraglutide or dapagliflozin. Similarly, no significant differences were observed in lean mass for any groups tested (Fig. 4B). Compared with HF controls, all treatments significantly decreased triglycerides ($p<0.5$ - $p<0.001$; Fig. 4C). DAPA-Lira reduced triglycerides (71-87% decrease; $p<0.01$ - $p<0.001$; Fig. 4C) to a greater extent than either liraglutide or dapagliflozin alone. No significant differences between HF groups in terms of total cholesterol was observed (Fig. 4D).

Effects of DAPA-Lira on terminal organ weight, hormones and biomarkers

Administration of DAPA-Lira and dapagliflozin resulted in a significant reduction in inguinal adipose weight ($p<0.05$; Fig. 5A). Liver and pancreatic weights were not significantly different in

any of the HF groups (Fig. 5B and 5C). Total plasma GLP-1 concentrations were significantly increased in all treatment groups ($p<0.001$) compared to HF controls with DAPA-Lira group displaying increased levels of total GLP-1 (12-18% increase; $p<0.01$ - $p<0.001$) compared to liraglutide or dapagliflozin alone (Fig. 5D). No significant differences were noted in ALT levels in HF mice (Fig. 5E). Both dapagliflozin and DAPA-Lira treatment groups displayed significantly elevated (1.0-fold; $p<0.01$ - $p<0.001$) plasma IL-6 levels compared to HF controls (Fig. 5F). Liraglutide only treated mice displayed reduced (48% decrease; $p<0.001$) IL-6 levels compared to DAPA-Lira treatment (Fig. 5F). All treatment groups resulted in significant reduction (33-43% decrease; $p<0.05$) in corticosterone concentrations compared to HF controls (Fig. 5G).

Effects of DAPA-Lira in novel object recognition task

During test trial, no significant difference was noted in the recognition index (RI) for the HF group indicating that they could not discriminate between novel and familiar object thereby exhibiting impaired cognition (Fig. 6A). All treatment groups and lean control group displayed significantly increased RI (1.1-1.3-fold; $p<0.01$ - $p<0.001$) when exposed to novel object compared to HF controls, thus highlighting preference to explore novel *versus* familiar object (Fig. 6B-6F). Open field assessment revealed no effect of any treatments on motor activity (speed and path length), anxiety (grooming events) and exploration (rearing events) (data not shown).

Effects of DAPA-Lira on islet morphology, pancreatic hormone content and mRNA gene expression

As shown in Fig. 7A, beta cell area was significantly increased ($p<0.05$) in liraglutide treated mice. Mice treated with dapagliflozin alone exhibited marked increase ($p<0.01$) in alpha cell area compared to HF controls (Fig. 7D). In contrast, DAPA-Lira or liraglutide alone did not affect alpha

cell area (Fig. 7D). Islets of HF mice exhibited substantial staining for IL-6 in beta cells and GLP-1 in alpha cells with no appreciable differences between various treatment groups (images not shown). Both DAPA-Lira and liraglutide markedly increased (1.4-1.6 fold; $p<0.05$) pancreatic insulin content compared to HF controls (Fig. 7B). Liraglutide treatment also significantly decreased (23% reduction; $p<0.001$) glucagon content while both DAPA-Lira and dapagliflozin led to significant increases compared to HF controls (21-28% increase; $p<0.01$ - $p<0.001$; Fig. 7E). A similar pattern to changes in pancreatic insulin and glucagon content were observed in pancreatic mRNA expression of insulin and glucagon (Fig. 7C and 7F).

Effects of DAPA-Lira on neurogenesis and synaptic density

Representative micrographs of doublecortin and synaptophysin staining are shown in Fig. 8A-8E. Mice treated with DAPA-Lira, dapagliflozin or liraglutide displayed increased number of immature neurons in the dentate gyrus (44-69% increase; $p<0.01$ - $p<0.001$; Fig. 8F) compared to HF controls as indicated by increased number of DCX-positive cells. Significantly higher levels of synaptophysin expression were demonstrated in all treatment groups in stratum oriens layer (88-113% increase; $p<0.01$; Fig. 8J) compared to HF controls. DAPA-Lira treatment also improved synaptophysin expression in stratum pyramidale layer (50% increase; $p<0.05$; Fig. 8I), though no differences in polymorph layer (Fig. 8G) and stratum radiatum (Fig. 8H) were observed.

Discussion

Given the increase and diversity of new antidiabetic drugs in the clinic, there is now a great opportunity to offer a more patient-centered tailored or personalized approach to therapeutic intervention. In this study, we examined the efficacy of combination therapy using the SGLT2 inhibitor dapagliflozin and the long-acting GLP-1 agonist liraglutide. In addition to assessing

metabolic outcomes, we also examined potential neuroprotective benefits of combination therapy on learning and memory, especially since recent evidence has shown that GLP-1 agonists may reduce cognitive decline in diabetes-obesity (Gault *et al.* 2010, Porter *et al.* 2013).

We chose to use a rodent model of diabetes combining low-dose STZ and high fat feeding to promote obesity, insulin resistance and hyperglycaemia (Srinivasan *et al.* 2005, Islam & Wilson 2012). This rodent model has been used previously and serves as a suitable means to evaluate potential drug intervention (Bhat *et al.* 2013, Millar *et al.* 2016). A small dose of STZ was combined with high fat feeding to inflict sub-lethal damage to beta cells which when combined with high fat feeding gave a more rapid and pronounced diabetes phenotype with elevation in blood glucose concentrations. SGLT2 inhibitors act by reabsorbing glucose so it is useful to study therapeutic effects when glucose levels are significantly raised. In contrast to liraglutide, DAPA-Lira combination therapy over 28 days did not affect energy intake. This is an important observation as several studies suggested that energy intake is increased following SGLT2 inhibitor therapy (Devenny *et al.* 2012, Nagata *et al.* 2013). As expected, all treatments resulted in reduced body weight, which in the case of dapagliflozin most presumably reflects energy loss via urinary glucose excretion (Scheen & Paquot 2014). SGLT2 inhibitors induce a significant and durable weight loss in patients with T2DM (Bailey *et al.* 2015).

Consistent with previous studies, monotherapy with dapagliflozin or liraglutide decreased glucose concentrations both in terms of non-fasting concentrations and following an oral glucose challenge. Moreover, DAPA-Lira combination therapy resulted in a more pronounced glucose-lowering effect that is most likely achieved through increased urinary glucose excretion (Bailey *et al.* 2016) and enhanced beta cell function. Interestingly, dapagliflozin monotherapy was also associated with enhanced glucose-induced insulin secretion and HOMA- β , which could be due to improved metabolic control and reversal of beta cell glucotoxicity, a potential direct effect on beta

cells and possible involvement of effects on other hormones such as GLP-1 as observed with less selective SGLT2 inhibitors (Zambrowicz *et al.* 2013). Indeed, plasma GLP-1 concentrations were significantly increased in all treatment groups at the end of the study but most notably in the group receiving DAPA-Lira combination therapy. This may well point to enhanced alpha cell GLP-1 production, which has been observed previously in pregnancy and situations of beta cell stress (Moffett *et al.* 2014, Vasu *et al.* 2014). Interestingly, IL-6 which has been implicated in islet processing of proglucagon to GLP-1 via increased expression of PC1/3 (Ellingsgaard *et al.* 2011) was markedly increased in beta cells of all HF groups.

All treatment groups exhibited improved insulin sensitivity and improved HOMA-IR. These changes in insulin sensitivity may be ascribed to weight reduction and alleviation of glucose toxicity (Macdonald *et al.* 2010). Of particular note is the observation that DAPA-Lira combination therapy markedly reduced HbA1c, which was significantly lower than either dapagliflozin or liraglutide alone. Importantly, no episodes of hypoglycaemia were observed in treatment groups following fasting for OGTT or during the insulin tolerance test, however measurement of circulating ketones would have been informative. Taken together, DAPA-Lira combination therapy was associated with improved glucose-lowering and greater reduction in body weight (compared to liraglutide alone), without observable effects on energy intake, suggesting that combination of dapagliflozin and liraglutide is a very powerful approach to management of glycaemia.

Consistent with previous studies, high fat fed mice exhibited dyslipidaemia and obesity (Podrini *et al.* 2013). DEXA scanning revealed that fat mass was significantly reduced in all treatment groups with tendency to be lower in groups treated with dapagliflozin and this was further corroborated by significantly decreased adipose tissue mass. In T2DM patients, body weight loss induced by dapagliflozin is mainly due to reduction in visceral and subcutaneous fat mass (Bolinder *et al.* 2012). Importantly, decreases in fat mass were not associated with changes in lean mass. High

fat mice displayed elevated triglyceride concentrations that were significantly improved in all treatment groups but especially in DAPA-Lira combination group. This could be due to function of improved glycaemia and greater weight reduction in this group. No significant changes in plasma total cholesterol were noted in any of the treatment groups. Although not measured in this study, relatively small clinically insignificant increases in both LDL- and HDL-cholesterol have been observed in patients on dapagliflozin therapy (Matthaei *et al.* 2015). As expected, HF mice exhibited increased liver weight and ALT concentrations characteristic of non-alcoholic fatty liver disease (Ganz *et al.* 2014). Whilst all treatments tended to reverse negative effects towards that of healthy controls, more detailed analyses investigating effects on hepatic triglyceride content, plasma and liver oxidative stress would be useful.

Mice treated with dapagliflozin, either alone or in combination with liraglutide, displayed elevated levels of IL-6. IL-6 has been shown to stimulate insulin from beta cells, glucagon from alpha cells and GLP-1 secretion from both intestinal L and pancreatic alpha cells (Ehse *et al.* 2007, Ellingsgaard *et al.* 2008, 2011). Increase in GLP-1 production is thought to occur via differentiation of the alpha cell through proglucagon transcription and enhanced PC1/3 expression (Ellingsgaard *et al.* 2008, 2011). Furthermore, SGLT2 inhibition promotes glucagon secretion from alpha cells in healthy mice (Bonner *et al.* 2015) and increases GLP-1 concentrations in patients (Ferrannini *et al.* 2014). More recent studies have suggested that dapagliflozin stimulates GLP-1 and IL6 secretion from pancreatic islets (Timper *et al.* 2016). In the present study, dapagliflozin increased plasma GLP-1, glucagon and IL-6, perhaps pointing to an important role for IL6 in beneficial action of dapagliflozin. Indeed, we have recently demonstrated increased expression of PC1/3 in α TC1.9 cells treated with dapagliflozin (unpublished observations). Measures of additional circulating cytokines (e.g. TNF α and IL-1 β) would provide insight as to whether this reflects a specific effect on IL-6, or a more generalized heightened inflammatory state. The increase in plasma IL-6 following

dapagliflozin therapy is particularly interesting and further studies exploring its role are clearly warranted. Importantly, mice treated with liraglutide alone displayed decreased plasma glucagon with no change in IL-6 concentrations.

Pancreatic immunochemical staining revealed that mice treated with dapagliflozin exhibited a significant increase in alpha cell area. This was accompanied by reduction in pancreatic insulin content and increases in both proglucagon gene expression and pancreatic glucagon content. These observations are broadly in line with metabolic insulin and glucagon data and further confirm an important effect of dapagliflozin on the alpha cell (Bonner *et al.* 2015). As expected, liraglutide treatment was associated with enhanced beta cell area and increased insulin content and this was accompanied by significant decrease in gene expression and pancreatic glucagon content (Schwasinger-Schmidt *et al.* 2013). Interestingly, alpha cell area was not affected in mice receiving DAPA-Lira combination therapy with both insulin gene expression and hormone content increased, suggesting that the liraglutide component countered some of the alpha cell promoting properties of dapagliflozin, which would be beneficial in a longer-term treatment regimen. We did not see tight correlation between the various parameters (percentage islet cells, hormone content and basal hormone levels) because many different factors influence these parameters. For example, if cell synthesizes hormone it is not just leaked out into the blood but is stored in vesicles that are regulated on minute-to-minute basis by prevailing blood glucose plus myriad of other factors.

We and others have previously shown that high fat feeding in rodents causes detrimental effects in brain regions associated with learning and memory (Greenwood & Winocur 2005, Stranahan *et al.* 2008a, Gault *et al.* 2010). Furthermore, growing evidence indicates that diabetes and obesity increase the risk of developing neurodegenerative disorders, such as Alzheimer's disease (Rani *et al.* 2016). More recently, GLP-1 agonists (and DPP4 inhibitors) have been shown to reduce cognitive decline associated with diabetes-obesity (Groeneveld *et al.* 2016). The present

study evaluated learning and memory using well-established novel object recognition task, which exploits ability of a rodent to explore a novel object over a familiar object (Abbas *et al.* 2009). As expected, HF mice could not discriminate between familiar and novel object when compared to healthy controls (Gault *et al.* 2015). However, all HF treated mice displayed significantly improved recognition memory, which was not attributable to effects on anxiety or motor activity, as indicated in open field assessment. Future studies using additional behavioural tests such as Morris Water Maze would also be useful.

Corticosterone concentrations were markedly raised following high fat feeding but treatment with dapagliflozin or liraglutide (both alone and in combination) reversed this effect. Raised glucocorticoid concentrations not only induce insulin resistance but also contribute to deficits in hippocampal function (Stranahan *et al.* 2008b). Indeed, reducing corticosterone concentrations can prevent diabetes-induced impairment of hippocampus-dependent learning (Stranahan *et al.* 2008c). Immunohistochemical staining revealed that HF treated mice displayed significantly enhanced doublecortin and synaptophysin expression indicating possible role of drug treatment to promote recovery of neurogenesis and synaptic density. It is possible that combination therapy over a longer time period may have resulted in superior neurogenesis and cognitive function compared to monotherapy. To our knowledge, this is the first study to report beneficial effects of dapagliflozin on cognitive function, neurogenesis, and synaptic density. Whilst positive neuroprotective and growth factor like effects of dapagliflozin on learning and memory are unlikely to occur as direct effect of SGLT2 inhibition in the brain itself, SGLT2 inhibitors are lipid-soluble and should cross the blood-brain-barrier (Bakris *et al.* 2009). However, similar to DPP4 inhibitors, it is plausible that neuroprotective effects observed for dapagliflozin could be attributed to increased GLP-1 concentrations which can then cross the blood-brain-barrier and/or actions of dapagliflozin to lower corticosterone concentrations. Whilst we cannot rule out that part of the neuroprotective effects may

be dependent on improved peripheral glycaemia, further detailed studies to delineate mechanism for this improvement in cognitive function with dapagliflozin would be useful.

In summary, compared with either agent alone, DAPA-Lira combination therapy was associated with superior glucose-lowering and significant reduction in body weight, indicating powerful and complementary approach to effectively manage hyperglycaemia. Part of this benefit appears to derive from the ability of liraglutide to decrease islet alpha cells. Other prominent effects included normalisation of hypertriglyceridaemia and enhancements of both insulin secretion and action. Furthermore, DAPA-Lira combination therapy, and indeed dapagliflozin monotherapy, were effective in reversing memory impairment in diabetic mice. Moreover, changes in glucagon and GLP-1 following dapagliflozin treatment were associated with changes in IL-6, suggesting possible role of IL-6 in mediating some of the actions of SGLT2 inhibition. This study supports recent papers showing clinical effectiveness of combination therapy with SGLT2 inhibition and stable GLP-1 mimetics in T2DM patients (DeFronzo 2017). Taken together, our results highlight an important personalised approach in utilising liraglutide in combination with dapagliflozin for further clinical evaluation in the treatment of diabetes and associated neurodegenerative disorders.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

These studies were supported by Department of Education and Learning PhD studentship to PJBM, Ulster University Strategic Research Funding, SAAD Trading and Contracting Company and

European Regional Development Fund (ERDF) award to AJB under the EU Sustainable Competitiveness Programme for Northern Ireland 2007-2013.

Author contributions

PJBM contributed to study design, conduct/data collection, data analysis and writing of the manuscript. VP, RCM, VParth and NMP contributed to conduct/data collection and data analysis. AJB and MO'K reviewed the manuscript. VAG and PRF contributed to study design, analysis and writing of the manuscript. All authors approved the final version of the manuscript.

References

Abbas T, Faivre E, Hølscher C, 2009 Impairment of synaptic plasticity and memory formation in GLP-1 receptor KO mice: interaction between type 2 diabetes and Alzheimer's disease. *Behavioural Brain Research* **205** 265-271.

Agersø H, Jensen LB, Elbrønd B, Rolan P, Zdravkovic M, 2002 The pharmacokinetics, pharmacodynamics, safety and tolerability of NN2211, a new long-acting GLP-1 derivative, in healthy men. *Diabetologia* **45** 195-202.

Ashraghi MR, Pagano G, Polychronis S, Niccolini F, Politis M, 2016 Parkinson's Disease, Diabetes and Cognitive Impairment. *Recent Patents on Endocrine, Metabolic and Immune Drug Discovery* **10** 11-21.

Bailey CJ, Morales Villegas EC, Woo V, Tang W, Ptaszynska A, List JF, 2015 Efficacy and safety of dapagliflozin monotherapy in people with type 2 diabetes: a randomized double-blind placebo-controlled 102-week trial. *Diabetic Medicine* **32** 531-541.

Bailey CJ, Tahrani AA & Barnett AH, 2016 Future glucose-lowering drugs for type 2 diabetes. *The Lancet Diabetes and Endocrinology* **4** 350-359.

Bakris GL, Fonseca VA, Skarma K, Wright EM, 2009 Renal sodium-glucose transport: role in diabetes mellitus and potential clinical implications. *Kidney International* **75** 1272-1277.

Bhat VK, Kerr BD, Flatt PR, Gault VA, 2013 A novel GIP-oxytomodulin hybrid peptide acting through GIP, glucagon and GLP-1 receptors exhibits weight reducing and anti-diabetic properties. *Biochemical Pharmacology* **85** 1655-1662.

Bolinder J, Ljunggren Ö, Kulberg J, Johansson L, Wilding J, Langkilde AM, Sugg J, Parikh S, 2012 Effects of dapagliflozin on body weight, total fat mass, and regional adipose tissue distribution in patients with type 2 diabetes mellitus with inadequate glycemic control on metformin. *Journal of Clinical Endocrinology and Metabolism* **97** 1020-1031.

Bonner C, Kerr-Conte J, Gmyr V, Queniat G, Moerman E, Thévenet J, Beaucamps C, Delalleau N, Popescu I, Malaisse WJ, *et al.* 2015 Inhibition of the glucose transporter SGLT2 with dapagliflozin in pancreatic alpha cells triggers glucagon secretion. *Nature Medicine* **21** 512-517.

da Rocha Fernandes J, Ogurtsova K, Linnenkamp U, Guariguata L, Seuring T, Zhang P, Cavan D, Makaroff LE, 2016 IDF Diabetes Atlas estimates of 2014 global health expenditures on diabetes. *Diabetes Research and Clinical Practice* **117** 48-54.

DeFronza RA, 2017 Combination therapy with GLP-1 receptor agonist and SGLT2 inhibitor. *Diabetes Obesity and Metabolism* [Epub ahead of print].

Devenny JJ, Godonis HE, Harvey SJ, Rooney S, Cullen MJ, Pelleymounter MA, 2012 Weight loss induced by chronic dapagliflozin treatment is attenuated by compensatory hyperphagia in diet-induced obese (DIO) rats. *Obesity (Silver Spring)* **20** 1645-1652.

Ehses JA, Perren A, Eppler E, Ribaux P, Pospisilik JA, Maor-Cahn R, Gueripel X, Ellingsgaard H, Schneider MK, Biollaz G, *et al.* 2007 Increased number of islet-associated macrophages in type 2 diabetes. *Diabetes* **56** 2356-2370.

Ellingsgaard H, Ehses JA, Hammar EB, Van Lommel L, Quintens R, Martens G, Kerr-Conte J, Pattou F, Berney T, Pipeleers D, *et al.* 2008 Interleukin-6 regulates pancreatic alpha-cells expansion. *Proceedings of the National Academy of Sciences of the United States of America* **105** 13163-13168.

Ellingsgaard H, Hauselmann I, Schuler B, Habib AM, Baggio LL, Meier DT, Eppler, E, Bouzakri K, Wueest S, Muller YD, *et al.* 2011 Interleukin-6 enhances insulin secretion by increasing glucagon-like peptide-1 secretion from L cells and alpha cells. *Nat Medicine* **17** 1481-1489.

- 524 Ferrannini E, Muscelli E, Frascerra S, Baldi S, Mari A, Heise T, Broedl UC, Woerle HJ, 2014
 525 Metabolic response to sodium-glucose cotransporter 2 inhibition in type 2 diabetic patients. *Journal*
 526 *of Clinical Investigation* **124** 499-508.
- 527
- 528 Flatt PR, Bailey CJ, 1981 Abnormal plasma glucose and insulin responses in heterozygous lean
 529 (ob/+) mice. *Diabetologia* **20** 573-577.
- 530
- 531 Ganz M, Csak T, Szabo G, 2014 High fat feeding results in gender specific steatohepatitis and
 532 inflammasome activation. *World Journal of Gastroenterology* **20** 8528-8534.
- 533
- 534 Gault VA, Porter WD, Flatt PR, Hölscher C, 2010 Actions of exendin-4 therapy on cognitive
 535 function and hippocampal synaptic plasticity in mice fed a high-fat diet. *International Journal of*
 536 *Obesity (London)* **34** 1341-1344.
- 537
- 538 Gault VA, Lennox R, Flatt PR, 2015 Sitagliptin, a dipeptidyl peptidase-4 inhibitor, improves
 539 recognition memory, oxidative stress and hippocampal neurogenesis and upregulates key genes
 540 involved in cognitive decline. *Diabetes Obesity and Metabolism* **17** 403-413.
- 541
- 542 Gerich JE, Meyer C, Woerle HJ, Stumvoll M, 2001 Renal gluconeogenesis: its importance in human
 543 glucose homeostasis. *Diabetes Care* **24** 382-391.
- 544
- 545 Greenwood CE, Winocur G, 2005 High-fat diets, insulin resistance and declining cognitive function.
 546 *Neurobiology of Aging* **26** 42-45.
- 547

Groeneveld ON, Kapelle LJ, Biessels GJ, 2016 Potentials of incretin-based therapies in dementia and stroke in type 2 diabetes mellitus. *Journal of Diabetes Investigation* **7** 5-16.

Han S, Hagan DL, Taylor JR, Xin L, Meng W, Biller SA, Wetterau JR, Washburn WN, Whaley JM, 2008 Dapagliflozin, a selective SGLT2 inhibitor, improves glucose homeostasis in normal and diabetic rats. *Diabetes* **57** 1723-1729.

Hediger MA, Rhoads DB, 1994 Molecular physiology of sodium-glucose cotransporters. *Physiological Reviews* **74** 993-1026.

Islam MS, Wilson RD, 2012 Experimentally induced rodent models of type 2 diabetes. *Methods in Molecular Biology* **933** 161-174.

Knudsen LB, Nielsen PF, Huusfeldt PO, Johansen NL, Madsen K, Pedersen FZ, Thøgersen H, Wilken M, Agersø H, 2000 Potent derivatives of glucagon-like peptide-1 with pharmacokinetic properties suitable for once daily administration. *Journal of Medicinal Chemistry* **43** 1664-1669.

Lennox R, Flatt PR, Gault VA, 2014 Lixisenatide improves recognition memory and exerts neuroprotective actions in high-fat fed mice. *Peptides* **61** 38-47.

Li Y, Zheng X, Meng F, Gong M, 2016 Application of self-assembling peptide as drug carrier for extending the GLP-1 stability. *International Journal of Clinical and Experimental Medicine* **9** 7828-7836.

572

573 Livak KJ, Schmittgen TD, 2001 Analysis of relative gene expression data using real-time
574 quantitative PCR and the 2(-Delta C(T)) Method. *Methods* **25** 402-408.

575

576 Macdonald FR, Peel JE, Jones HB, Mayers RM, Westgate L, Whaley JM, Poucher SM, 2010 The
577 novel sodium glucose transporter 2 inhibitor dapagliflozin sustains pancreatic function and preserves
578 islet morphology in obese, diabetic rats. *Diabetes Obesity and Metabolism* **12** 1004-1012.

579

580 Madsen K, Knudsen LB, Agersø H, Nielsen PF, Thøgersen H, Wilken M, Johansen NL, 2007
581 Structure-activity and protraction relationship of long-acting glucagon-like peptide-1 derivatives:
582 importance of fatty acid length, polarity, and bulkiness. *Journal of Medicinal Chemistry* **50** 6126-
583 6132.

584

585 Matthaei S, Bowering K, Rohwedder K, Sugg J, Parikh S, Johnsson E, 2015 Study 05 Group.
586 Durability and tolerability of dapagliflozin over 52 weeks as add-on to metformin and sulphonylurea
587 in type 2 diabetes. *Diabetes Obesity and Metabolism* **17** 1075-1084.

588

589 McClean PL, Parthasarathy V, Faivre E, Hölscher C, 2011 The diabetes drug liraglutide prevents
590 degenerative processes in a mouse model of Alzheimer's disease. *Journal of*
591 *Neuroscience* **31** 6587-6594.

592

593 Merovci A, Mari A, Solis C, Xiong J, Daniele G, Chavez-Velazquez A, Tripathy D, Urban
594 McCarthy S, Abdul-Ghani M, DeFronzo RA, 2015 Dapagliflozin lowers plasma glucose

concentration and improves beta-cell function. *Journal of Clinical Endocrinology and Metabolism* **100** 1927-1932.

Millar PJ, Pathak V, Moffett RC, Pathak NM, Bjourson AJ, O’Kane MJ, Flatt PR, Gault VA, 2016 Beneficial metabolic actions of a stable GIP agonist following pre-treatment with a SGLT2 inhibitor in high fat fed diabetic mice. *Molecular and Cellular Endocrinology* **420** 37-45.

Moffett RC, Vasu S, Thorens B, Drucker DJ, Flatt PR, 2014 Incretin receptor null mice reveal key role of GLP-1 but not GIP in pancreatic beta cell adaptation to pregnancy. *PLoS One* 9(6):e96863

Moffett RC, Patterson S, Irwin N, Flatt PR, 2015 Positive effects of GLP-1 receptor activation with liraglutide on pancreatic islet morphology and metabolic control in C57BL/KsJ db/db mice with degenerative diabetes. *Diabetes Metabolism Research and Reviews* **31** 248-255.

Mudaliar S, Henry RR, Boden G, Smith S, Chalamandaris AG, Duchesne D, Iqbal N, List J, 2014 Changes in insulin sensitivity and insulin secretion with the sodium glucose cotransporter 2 inhibitor dapagliflozin. *Diabetes Technology and Therapeutics* **16** 137-144.

Nagata T, Fukuzawa T, Takeda M, Fukazawa M, Mori T, Nihei T, Honda K, Suzuki Y, Kawabe Y, 2013 Tofogliflozin, a novel sodium-glucose co-transporter 2 inhibitor, improves renal and pancreatic function in db/db mice. *British Journal of Pharmacology* **170** 519-531.

Parthsarathy V, McClean PL, Hölscher C, Taylor M, Tinker C, Jones G, Kolosov O, Salvati E, Gregori M, Masserini M, Allsop D, 2013 A Novel Retro-Inverso Peptide Inhibitor Reduces

Amyloid Deposition, Oxidation and Inflammation and Stimulates Neurogenesis in the APPswe/PS1ΔE9 Mouse Model of Alzheimer's Disease. *PLoS One* **8**(1): e54769.

Podrini C, Cambridge EL, Lelliott CJ, Carragher DM, Estabel J, Gerdin AK, Karp NA, Scudamore CL; Sanger Mouse Genetics Project., Ramirez-Solis R, White JK, 2013 High-fat feeding rapidly induces obesity and lipid derangements in C57BL/6N mice. *Mammalian Genome* **24** 240-251.

Porter WD, Flatt PR, Hölscher C, Gault VA, 2013 Liraglutide improves hippocampal synaptic plasticity associated with increased expression of Mash1 in ob/ob mice. *International Journal of Obesity (London)* **37** 678-684.

Rahmoune H, Thompson PW, Ward JM, Smith CD, Hong G, Brown J, 2005 Glucose transporters in human renal proximal tubular cells isolated from the urine of patients with non-insulin-dependent diabetes. *Diabetes* **54** 3427-3434.

Rani V, Deshmukh R, Jaswal P, Kumar P, Bariwal J, 2016 Alzheimer's disease: is this a brain specific diabetic condition? *Physiology and Behavior* **164** 259-267.

Renner S, Blutke A, Streckel E, Wanke R, Wolf E, 2016 Incretin actions and consequences of incretin-based therapies: lessons from complementary animal models. *The Journal of Pathology* **238** 345-358.

Scheen AJ, Paquot N, 2014 Metabolic effects of SGLT2 inhibitors beyond increased glucosuria: a review of the clinical evidence. *Diabetes and Metabolism* **40** S4-S11.

Schwasinger-Schmidt T, Robbins DC, Williams SJ, Noyikova L, Stehno-Bittel L, 2013 Long-term liraglutide treatment is associated with increased insulin content and secretion in beta cells, and a loss of alpha cells in ZDF rats. *Pharmacological Research* **76** 58-66.

Srinivasan K, Viswanad B, Asrat L, Ramarao KP, 2005 Combination of high-fat diet-fed and low-dose streptozotocin-treated rat: a model for type 2 diabetes and pharmacological screening. *Pharmacological Research* **52** 313-320.

Stranahan AM, Norman ED, Lee K, Cutler RG, Telljohann RS, Egan JM, Mattson MP, 2008a Diet-induced insulin resistance impairs hippocampal synaptic plasticity and cognition in middle-aged rats. *Hippocampus* **18** 1085-1088.

Stranahan AM, Arumugam TV, Cutler RG, Lee K, Egan JM, Mattson MP, 2008b Diabetes impairs hippocampal function through glucocorticoid-mediated effects on new and mature neurons. *Nature Neuroscience* **11** 309-317.

Stranahan AM, Lee K, Pistell PJ, Nelson CM, Readal N, Miller MG, Spangler EL, Ingram DK, Mattson MP, 2008c Accelerated cognitive aging in diabetic rats is prevented by lowering corticosterone levels. *Neurobiology of Learning and Memory* **90** 479-483.

Timper K, Dalmás E, Dror E, Rütts, Thienel C, Sauter NS, Bouzakri K, Bédar B, Pattou F, Kerr-Conte J, 2016 Glucose-dependent insulintropic peptide stimulates glucagon-like peptide 1 production by pancreatic islets via interleukin 6, produced by α cells. *Gastroenterology* **151** 165-179.

668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691

Tramutola A, Arena A, Cini C, Butterfield DA, Barone E, 2017 Modulation of GLP-1 signaling as a novel therapeutic approach in the treatment of Alzheimer’s disease pathology. *Expert Review of Neurotherapeutics* **17** 59-75.

Vasu S, Moffett RC, Thorens B, Flatt PR, 2014 Role of endogenous GLP-1 and GIP in beta cell compensatory responses to insulin resistance and cellular stress. *PLoS One* **9**(6):e101005

Vivian EM, 2015 Dapagliflozin: a new sodium-glucose cotransporter 2 inhibitor for treatment of type 2 diabetes. *American Journal of Health-System Pharmacy* **72** 361-372.

Zaccardi F, Webb DR, Yates T, Davies MJ, 2016 Pathophysiology of type 1 and type 2 diabetes mellitus: a 90-year perspective. *Postgraduate Medical Journal* **92** 63-69.

Zambrowicz B, Ding ZM, Ogbaa I, Frazier K, Banks P, Turnage A, Freiman J, Smith M, Ruff D, Sands A, Powell D, 2013 Effects of LX4211, a dual SGLT1/SGLT2 inhibitor, plus sitagliptin on postprandial active GLP-1 and glycaemic control in type 2 diabetes. *Clinical Therapeutics* **35** 273-285.

Figure legends

Figure 1: Effects of once-daily administration of DAPA-Lira on (A) body weight, (B) cumulative energy intake, (C) glucose and (D) insulin concentrations. HF mice received saline vehicle (0.9% wt/vol; p.o.), dapagliflozin (1 mg/kg; p.o.), dapagliflozin (1 mg/kg; p.o.) plus liraglutide (25 nmol/kg; i.p) or liraglutide alone (25 nmol/kg; i.p) over 28 days. Lean control mice received saline vehicle once-daily. Metabolic parameters were measured every 3 to 4 days. Values are means \pm SEM for groups of 8-10 mice. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ compared to HF controls. $^{\Delta\Delta}p < 0.01$ compared to DAPA-Lira.

Figure 2: Effects of once-daily administration of DAPA-Lira on (A and B) glucose tolerance and (C and D) insulin response to glucose. HF mice received saline vehicle (0.9% wt/vol; p.o.), dapagliflozin (1 mg/kg; p.o.), dapagliflozin (1 mg/kg; p.o.) plus liraglutide (25 nmol/kg; i.p) or liraglutide alone (25 nmol/kg; i.p) over 28 days. Lean control mice received saline vehicle once-daily. Glucose and insulin concentrations were measured prior to and after oral administration of glucose (18 mmol/kg) in 12-hour fasted mice. Glucose and insulin AUC values post-injection are also shown. Values are means \pm SEM for groups of 8-10 mice. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ compared to HF controls. $^{\Delta\Delta}p < 0.01$ and $^{\Delta\Delta\Delta}p < 0.001$ compared to DAPA-Lira.

Figure 3: Effects of once-daily administration of DAPA-Lira on (A and B) insulin sensitivity, (C) HOMA-IR, (D) HOMA- β , (E) HbA1c and (F) plasma glucagon. HF mice received saline vehicle (0.9% wt/vol; p.o.), dapagliflozin (1 mg/kg; p.o.), dapagliflozin (1 mg/kg; p.o.) plus liraglutide (25 nmol/kg; i.p) or liraglutide alone (25 nmol/kg; i.p) over 28 days. Lean control mice received saline vehicle once-daily. Parameters were measured at the end of the study period. For insulin sensitivity, glucose concentrations were measured prior to and after injection of insulin (25 U/kg; i.p.) in non-

fasted mice. Glucose AAC values post-injection are also shown. Values are means \pm SEM for groups of 8-10 mice. $**p<0.01$ and $***p<0.001$ compared to HF controls. $^{\Delta}p<0.05$, $^{\Delta\Delta}p<0.01$ and $^{\Delta\Delta\Delta}p<0.001$ compared to DAPA-Lira.

Figure 4: Effects of once-daily administration of DAPA-Lira on (A) fat mass, (B) lean mass, (C) triglycerides and (D) total-cholesterol. HF mice received saline vehicle (0.9% wt/vol; p.o.), dapagliflozin (1 mg/kg; p.o.), dapagliflozin (1 mg/kg; p.o.) plus liraglutide (25 nmol/kg; i.p) or liraglutide alone (25 nmol/kg; i.p) over 28 days. Lean control mice received saline vehicle once-daily. Parameters were measured at the end of the study period. Values are means \pm SEM for groups of 8-10 mice. $*p<0.05$ and $***p<0.001$ compared to HF controls. $^{\Delta\Delta}p<0.01$ and $^{\Delta\Delta\Delta}p<0.001$ compared to DAPA-Lira.

Figure 5: Effects of once-daily administration of DAPA-Lira on (A) adipose weight, (B) liver weight, (C) pancreatic weight, (D) total GLP-1, (E) ALT, (F) corticosterone, and (G) IL-6 concentrations. HF mice received saline vehicle (0.9% wt/vol; p.o.), dapagliflozin (1 mg/kg; p.o.), dapagliflozin (1 mg/kg; p.o.) plus liraglutide (25 nmol/kg; i.p) or liraglutide alone (25 nmol/kg; i.p) over 28 days. Lean control mice received saline vehicle once-daily. Parameters were measured at the end of the study period. Values are means \pm SEM for groups of 8-10 mice. $*p<0.05$, $**p<0.01$ and $***p<0.001$ compared to HF controls. $^{\Delta\Delta}p<0.01$ and $^{\Delta\Delta\Delta}p<0.001$ compared to DAPA-Lira.


Figure 6: Effects of once-daily administration of DAPA-Lira on recognition index for (A) HF, (B) DAPA, (C) DAPA-Lira, (D) Lira, (E) lean mice and (F) difference score. HF mice received saline vehicle (0.9% wt/vol; p.o.), dapagliflozin (1 mg/kg; p.o.), dapagliflozin (1 mg/kg; p.o.) plus liraglutide (25 nmol/kg; i.p) or liraglutide alone (25 nmol/kg; i.p) over 28 days. Lean control mice

received saline vehicle once-daily. Parameters were measured at the end of the study period. Values are means \pm SEM for groups of 8-10 mice. $*p<0.05$, $**p<0.01$ and $***p<0.001$ compared to HF controls. $^{\Delta}p<0.05$ compared to DAPA-Lira.

Figure 7: Effects of once-daily administration of DAPA-Lira on (A) beta cell area, (B) insulin content, (C) insulin mRNA expression, (D) alpha cell area, (E) glucagon content and (F) glucagon mRNA expression. HF mice received saline vehicle (0.9% wt/vol; p.o.), dapagliflozin (1 mg/kg; p.o.), dapagliflozin (1 mg/kg; p.o.) plus liraglutide (25 nmol/kg; i.p) or liraglutide alone (25 nmol/kg; i.p) over 28 days. Lean control mice received saline vehicle once-daily. Parameters were measured at the end of the study period. Values are means \pm SEM for groups of 8-10 mice. $*p<0.05$, $**p<0.01$ and $***p<0.001$ compared to HF controls. $^{\Delta\Delta}p<0.01$ and $^{\Delta\Delta\Delta}p<0.001$ compared to DAPA-Lira.


Figure 8: Effects of once-daily administration of DAPA-Lira on (A-E) brain immunohistochemistry, (F) doublecortin neuroblast, and quantification levels of synaptophysin expression in (G) polymorph layer, (H) stratum radiatum, (I) stratum pyramidale and (J) stratum oriens layer. HF mice received saline vehicle (0.9% wt/vol; p.o.), dapagliflozin (1 mg/kg; p.o.), dapagliflozin (1 mg/kg; p.o.) plus liraglutide (25 nmol/kg; i.p) or liraglutide alone (25 nmol/kg; i.p) over 28 days. Lean control mice received saline vehicle once-daily. Parameters were measured at the end of the study period. Values are means \pm SEM for groups of 6 mice. $*p<0.05$, $**p<0.01$ and $***p<0.001$ compared to HF controls.

Figure 1




254x190mm (96 x 96 DPI)

Figure 2




254x190mm (96 x 96 DPI)

Figure 3




254x190mm (96 x 96 DPI)

Figure 4




254x190mm (96 x 96 DPI)

Figure 5




254x190mm (96 x 96 DPI)

Figure 6




254x190mm (96 x 96 DPI)

Figure 7



254x190mm (96 x 96 DPI)

Figure 8



254x190mm (96 x 96 DPI)